

Closed-form two-loop Euler-Heisenberg Lagrangian in a self-dual background

Gerald V. Dunne

*Department of Physics
University of Connecticut
Storrs CT 06269, USA*

Christian Schubert

*Instituto de Física y Matemáticas
Universidad Michoacana de San Nicolás de Hidalgo
Apdo. Postal 2-82
C.P. 58040, Morelia, Michoacán, México
schubert@itzel.ifm.umich.mx*

and

*California Institute for Physics and Astrophysics
366 Cambridge Ave., Palo Alto, CA 94306, USA*

Abstract

We show that the two-loop Euler-Heisenberg effective Lagrangian for scalar QED in a constant Euclidean self-dual background has a simple explicit closed form expression in terms of the digamma function. This result leads to a simple analysis of the weak- and strong-field expansions, the two-loop scalar QED beta function, and the analytic continuation properties of the effective Lagrangian and its imaginary part.

The Euler-Heisenberg Lagrangian, one of the earliest results in quantum electrodynamics [1,2], describes the complete one-loop amplitude involving a spinor loop interacting non-perturbatively with a constant background field. This effective Lagrangian encodes the information concerning the one-loop amplitude in a form which is extremely convenient for the study at low energies of nonlinear QED effects such as photon-photon scattering [1], photon dispersion [3], and photon splitting [3]. Analogous results apply for scalar QED [4]. The Euler-Heisenberg Lagrangian is real for a purely magnetic field, while in the presence of an electric field there is an absorptive part, indicative of the possibility of electron-positron pair creation by the field [4,5]. Although pair production is exponentially suppressed for field strengths which are presently possible for a macroscopic field in the laboratory, pair creation has recently been observed in an experiment involving electrons traversing the focus of a terawatt laser [6]. Moreover, in the near future the construction of X-ray free electron lasers may allow one to obtain field strengths much higher than the ones obtainable by optical lasers [7].

All these results are at the one-loop level. Going to the two-loop level, the first radiative correction to the Euler-Heisenberg Lagrangian, describing the effect of an additional photon exchange in the loop, was first studied in the seventies by Ritus [8–10], for both spinor and scalar QED, and for a general constant background field $F_{\mu\nu}$. Using the exact propagators in a constant field found by Fock [11] and Schwinger [4], and a proper-time cutoff as the UV regulator, Ritus obtained the two-loop effective Lagrangian in terms of a certain two-parameter integral. A new feature of the two-loop calculation is the necessity of mass renormalization, which turns out to work in a quite non-trivial way [8–10,12–15]. Unfortunately, the double parameter integral is very complicated, so it is much more difficult to study the weak- and strong-field expansions at two-loops than at one-loop. This is true even for the special cases where the background is purely magnetic or purely electric [8–10]. More recently, the world-line approach to quantum field theory [16–21] has been used [13,14,22] to re-compute these two-loop results, for both spinor and scalar QED, in a general constant background field. The world-line results are once again expressed as complicated double parameter integrals, although they are of a different form from Ritus’s expressions. While it has not yet been shown how to convert between these two different expressions, their equivalence has been checked, as weak- and strong-field expansions, using computer-based expansions. The Borel summation properties of the weak-field expansion have also been studied [15].

In this letter we specialize the world-line two-loop calculation to a constant Euclidean self-dual background, where $F_{\mu\nu} = \tilde{F}_{\mu\nu} \equiv \frac{1}{2}\varepsilon_{\mu\nu\alpha\beta}F^{\alpha\beta}$. In this case we find the surprising result that *both* of the parameter integrals can be done in closed form, yielding a simple expression (see Eq.[18]) for the scalar QED effective Lagrangian, in terms of the digamma function $\psi(x) = d/dx \ln \Gamma(x)$. This simple closed form makes it extremely easy to study the weak- and strong-field expansions, as well as analyticity features such as analytic continuation and Borel summation. We have chosen to consider scalar QED rather than spinor QED as it leads to some computational simplification in the world-line formalism, but analogous simplifications occur for spinor QED [23].

A constant Euclidean self-dual background is of interest for a number of reasons. First, such backgrounds are significant in QCD, since constant gauge fields are only stable under quantum fluctuations if they are self-dual and essentially abelian (*i.e.*, have a fixed direction in color space) [24]. Second, a constant self-dual background is relevant for the derivative expansion for an instanton background in a non-abelian theory [25]. Third, this type of background plays an interesting role in the interpretation of D-instantons and D3-branes in $\mathcal{N} = 4$ super-Yang-Mills theory [26,27]. Fourth, interesting properties of the $SU(2)$ effective potential have been found recently for such self-dual backgrounds [28]. Finally, from a computational perspective, the self-dual case is the simplest nontrivial constant field background, and we will see that it leads to the most explicit expressions for the effective Lagrangian; hence we hope that it may lead to progress concerning the general properties of radiative corrections to the Euler-Heisenberg Lagrangian.

We begin by briefly recalling the one-loop results for scalar QED. The one-loop Euler-Heisenberg effective Lagrangian for scalar QED in a general constant background field $F_{\mu\nu}$ can be expressed as [4]

$$\mathcal{L}_{\text{scalar}}^{(1)} = \frac{1}{(4\pi)^2} \int_0^\infty \frac{dT}{T} e^{-m^2 T} \left[\frac{e^2 a b}{\sinh(eaT) \sin(ebT)} - \frac{1}{T^2} + \frac{e^2(a^2 - b^2)}{6} \right] \quad (1)$$

Here T denotes the (Euclidean) proper-time of the loop fermion, and a, b are related to the two Lorentz invariants of the Maxwell field by $a^2 - b^2 = \mathbf{B}^2 - \mathbf{E}^2$, and $ab = \mathbf{E} \cdot \mathbf{B}$. The last term in the square brackets in (1) corresponds to the one-loop charge renormalization [4].

If the field has an electric component then this effective Lagrangian has an imaginary part, indicating that the vacuum becomes unstable in an electric field. A convenient representation for this imaginary part was found by Schwinger [4]. In the case of a purely electric field E it reads

$$\text{Im}\mathcal{L}_{\text{scalar}}^{(1)}(E) = -\frac{e^2 E^2}{16\pi^3} \sum_{k=1}^{\infty} \frac{(-1)^k}{k^2} \exp\left[-\frac{m^2 \pi k}{eE}\right] \quad (2)$$

In this representation the coefficient of the k -th exponential can be directly identified with the rate for the production of k coherent pairs by the field [4,5].

For a constant Euclidean self-dual (SD) background field one can write $F^2 = -f^2 \mathbf{1}$, so that $a = f$, and $b = -if$. Then, specializing (1) to such a background,

$$\mathcal{L}_{\text{scalar}}^{(1)(SD)}(\kappa) = \frac{m^4}{(4\pi)^2} \frac{1}{4\kappa^2} \int_0^\infty \frac{dt}{t^3} e^{-2\kappa t} \left[\frac{t^2}{\sinh^2(t)} - 1 + \frac{t^2}{3} \right] \quad (3)$$

where we have defined the dimensionless parameter

$$\kappa \equiv \frac{m^2}{2e\sqrt{f^2}} \quad (4)$$

and rescaled $t = Te\sqrt{f^2}$.

The one-loop effective Lagrangian in (3) may be expressed in terms of special functions (such as zeta functions) in various ways [29]. However, in anticipation of a comparison with our two-loop result below, we present the following representation in terms of the function $\ln \Gamma(x)$ and its integral:

$$\mathcal{L}_{\text{scalar}}^{(1)(SD)}(\kappa) = \frac{m^4}{(4\pi)^2} \frac{1}{\kappa^2} \left[-\frac{1}{12} \ln \kappa + \zeta'(-1) + \Xi(\kappa) \right] \quad (5)$$

Here we have defined the function [30]

$$\Xi(x) \equiv -x \ln \Gamma(x) + \frac{x^2}{2} \ln x - \frac{x^2}{4} - \frac{x}{2} + \int_0^x dy \ln \Gamma(y) \quad (6)$$

and $\zeta'(-1) \approx -0.16542$. It is straightforward to deduce the weak- and strong-field expansions of the one-loop effective Lagrangian $\mathcal{L}_{\text{scalar}}^{(1)(SD)}(\kappa)$. The weak field (*i.e.*, large κ) expansion follows most easily from the integral representation (3):

$$\mathcal{L}_{\text{scalar}}^{(1)(SD)}(\kappa) = \frac{m^4}{(4\pi)^2} \sum_{n=2}^{\infty} c_n^{(1)} \frac{1}{\kappa^{2n}} \quad (7)$$

where the expansion coefficients are (here $\mathcal{B}_n$ are Bernoulli numbers)

$$c_n^{(1)} = -\frac{\mathcal{B}_{2n}}{2n(2n-2)} \quad , \quad n = 2, 3, \dots \quad (8)$$

The weak-field expansion (7) is a divergent alternating series, since the leading behavior of $c_n^{(1)}$ at large order is

$$c_n^{(1)} \sim 2 \frac{(-1)^n}{(2\pi)^{2n}} \Gamma(2n-1) \quad , \quad n \rightarrow \infty \quad (9)$$

The strong-field (*i.e.*, small κ) expansion of $\mathcal{L}_{\text{scalar}}^{(1)(SD)}(\kappa)$ follows most easily from the representation (5):

$$\mathcal{L}_{\text{scalar}}^{(1)(SD)}(\kappa) = \frac{m^4}{(4\pi)^2} \frac{1}{\kappa^2} \left[\left(-\frac{1}{12} + \frac{\kappa^2}{2} \right) \ln \kappa + \zeta'(-1) + \frac{\kappa}{2} + \left(\frac{\gamma}{2} - \frac{1}{4} \right) \kappa^2 - \sum_{n=2}^{\infty} \frac{(-1)^n \zeta(n)}{(n+1)} \kappa^{n+1} \right] \quad (10)$$

The leading strong-field term is

$$\mathcal{L}_{\text{scalar}}^{(1)(SD)} \sim \frac{1}{12\pi} \frac{e^2}{4\pi} f^2 \ln \left(\frac{2ef}{m^2} \right) \quad (11)$$

The scalar QED β -function at one-loop can be read off from the coefficient of the $f^2 \ln f$ term [8–10,31]. In the conventions of [32],

$$\beta_{\text{scalar}}(\alpha) = \frac{\alpha^2}{6\pi} + \dots \quad (12)$$

where $\alpha = \frac{e^2}{4\pi}$ is the fine structure constant. This illustrates the connection between the strong field and high momentum limits of scalar QED [8–10,31].

Turning to the two-loop case, we briefly recall the world-line results [13,14,22] for the two-loop Euler-Heisenberg Lagrangian in a general constant field $F_{\mu\nu}$:

$$\begin{aligned} \mathcal{L}_{\text{scalar}}^{(2)}(F) &= (4\pi)^{-D} \left(-\frac{e^2}{2} \right) \int_0^\infty \frac{dT}{T} e^{-m^2 T} T^{-\frac{D}{2}} \int_0^\infty d\bar{T} \int_0^T d\tau_a \int_0^T d\tau_b \\ &\quad \times \det^{-\frac{1}{2}} \left[\frac{\sin(eFT)}{eFT} \right] \det^{-\frac{1}{2}} \left[\bar{T} - \frac{1}{2} \mathcal{C}_{ab} \right] \langle \dot{y}_a \cdot \dot{y}_b \rangle \end{aligned} \quad (13)$$

Here T and $\bar{T}$ represent the scalar and photon proper-times, and $\tau_{a,b}$ the endpoints of the photon insertion moving around the scalar loop. $\mathcal{C}_{ab}$ and $\langle \dot{y}_a \cdot \dot{y}_b \rangle$ are given by

$$\begin{aligned} \mathcal{C}_{ab} &= \mathcal{G}_B(\tau_a, \tau_a) - \mathcal{G}_B(\tau_a, \tau_b) - \mathcal{G}_B(\tau_b, \tau_a) + \mathcal{G}_B(\tau_b, \tau_b) \\ \langle \dot{y}_a \cdot \dot{y}_b \rangle &= \text{tr} \left[\ddot{\mathcal{G}}_{Bab} + \frac{1}{2} \frac{(\dot{\mathcal{G}}_{Baa} - \dot{\mathcal{G}}_{Bab})(\dot{\mathcal{G}}_{Bab} - \dot{\mathcal{G}}_{Bbb})}{\bar{T} - \frac{1}{2} \mathcal{C}_{ab}} \right] \end{aligned} \quad (14)$$

They are expressed in terms of the worldline Green's function $\mathcal{G}_B$ (and its first and second derivatives), where [13]:

$$\mathcal{G}_B(\tau_1, \tau_2) = \frac{1}{2(eF)^2} \left(\frac{eF}{\sin(eFT)} e^{-ieFT\dot{\mathcal{G}}_{B12}} + ieF\dot{\mathcal{G}}_{B12} - \frac{1}{T} \right) \quad (15)$$

Here $\dot{\mathcal{G}}_{B12} = \text{sign}(\tau_1 - \tau_2) - 2(\tau_1 - \tau_2)/T$, and a ‘dot’ always refers to a derivative with respect to the first variable. The formula (15) should be understood as a power series in the field strength tensor $F_{\mu\nu}$.

We now specialize these two-loop formulas to the case of a self-dual field. The worldline correlator (15) for such a field simplifies to [33]:

$$\mathcal{G}_{B12} = \frac{T}{2} \left[\frac{1}{Z^2} - \frac{\cosh(Z\dot{\mathcal{G}}_{B12})}{Z \sinh(Z)} \right] \mathbb{1} + i \frac{T}{2Z^2} \left[\frac{\sinh(Z\dot{\mathcal{G}}_{B12})}{\sinh(Z)} - \dot{\mathcal{G}}_{B12} \right] \mathcal{Z} \quad (16)$$

where $\mathcal{Z}_{\mu\nu} \equiv eTF_{\mu\nu}$ and $Z \equiv eFT$. Contrary to the magnetic or electric field cases, the resulting integrand in (13) is simple enough to allow one to perform, in closed form, both the integrals over $\bar{T}$ and over $\tau_1 - \tau_2$. The result reads, after a partial integration in T ,

$$\mathcal{L}_{\text{scalar}}^{(2)(SD)}(\kappa) = \frac{m^4}{(4\pi)^4} \frac{e^2}{4\kappa^2} \int_0^\infty \frac{dt}{t^3} e^{-2\kappa t} \left(\frac{t}{\sinh(t)} \right)^2 \left\{ -4t^2 + 6\kappa t \left[1 - \gamma + \ln \left(\frac{1}{2\kappa \sinh(t)} \right) \right] \right\} \quad (17)$$

Here we have subtracted the appropriate mass renormalization term, as explained in [14]. The (on-shell) renormalization of the Lagrangian is completed by further subtracting from the integrand the terms of order $O(f^0), O(f^2)$. The resulting finite proper-time integral in (17) can then again be performed by elementary means, leading to the following simple expression for the renormalized two-loop effective Lagrangian:

$$\mathcal{L}_{\text{scalar}}^{(2)(SD)}(\kappa) = \frac{m^4 \alpha}{(4\pi)^3} \frac{1}{\kappa^2} \left[\frac{3}{2} \xi^2(\kappa) - \xi'(\kappa) \right] \quad (18)$$

where we define [here $\psi(x)$ is the digamma function: $\psi(x) = \frac{d}{dx} \ln \Gamma(x)$]

$$\xi(x) \equiv -x \left(\psi(x) - \ln x + \frac{1}{2x} \right) \quad (19)$$

The simplicity of our result (18) is very surprising, and we do not know of any comparable two-loop result in gauge theory (for ϕ^4 theory an analogous result was obtained in [34]). Comparing with the one-loop self-dual result (5), we note that the two-loop result (18) is very similar, since $\xi(x) = \Xi'(x)$. In fact, in some sense the two-loop result is even simpler than the one-loop result, as it involves derivatives of $\ln \Gamma(x)$ rather than its integral.

It is now straightforward to derive the weak- and strong-field expansions of the two-loop effective Lagrangian, using the known expansions of the digamma function:

$$\psi(x) = \begin{cases} \ln x - \frac{1}{2x} - \sum_{n=1}^{\infty} \frac{B_{2n}}{2nx^{2n}} & , \quad x \rightarrow \infty \\ -\frac{1}{x} - \gamma + \sum_{n=2}^{\infty} (-1)^n \zeta(n) x^{n-1} & , \quad |x| < 1 \end{cases} \quad (20)$$

Therefore, the weak-field (*i.e.*, large κ) expansion is

$$\mathcal{L}_{\text{scalar}}^{(2)(SD)}(\kappa) = \frac{m^4}{(4\pi)^2} \alpha \pi \sum_{n=2}^{\infty} c_n^{(2)} \frac{1}{\kappa^{2n}} \quad (21)$$

with two-loop expansion coefficients (for $n = 2, 3, \dots$):

$$c_n^{(2)} = \frac{1}{(2\pi)^2} \left\{ \frac{2n-3}{2n-2} B_{2n-2} + \frac{3}{2} \sum_{k=1}^{n-1} \frac{B_{2k}}{2k} \frac{B_{2n-2k}}{(2n-2k)} \right\} \quad (22)$$

At large order in the weak field expansion the leading behavior is

$$c_n^{(2)} \sim 2 \frac{(-1)^n}{(2\pi)^{2n}} \Gamma(2n-1) \quad , \quad n \rightarrow \infty \quad (23)$$

Thus, the leading growth rate (at large order of perturbation theory) of the weak-field expansion coefficients is exactly the same at two-loop as at one-loop, up to an overall factor of $\alpha\pi$. This is consistent with the results in [8–10,15] for the purely magnetic and electric cases. We have also checked (to 24 orders) that the weak-field expansion coefficients in (22) agree with those obtained by an expansion of Ritus's double-parameter integral expression in [9].

Similarly, from (18,19,20), the strong-field (*i.e.*, small κ) expansion is

$$\begin{aligned} \mathcal{L}_{\text{scalar}}^{(2)(SD)}(\kappa) = \frac{m^4 \alpha}{(4\pi)^3} \frac{1}{\kappa^2} & \left[(\gamma + \ln \kappa) \left\{ -1 + \frac{3}{2} \kappa + \frac{3}{2} \kappa^2 (\gamma + \ln \kappa) - 3 \sum_{n=2}^{\infty} (-1)^n \zeta(n) \kappa^{n+1} \right\} - \frac{5}{8} + \frac{\pi^2}{3} \kappa \right. \\ & \left. - \sum_{n=2}^{\infty} (-1)^n \left(\frac{3}{2} \zeta(n) + (n+1) \zeta(n+1) \right) \kappa^n + \frac{3}{2} \sum_{n=4}^{\infty} \sum_{l=2}^{n-2} (-1)^n \zeta(n-l) \zeta(l) \kappa^n \right] \end{aligned} \quad (24)$$

The leading strong-field term is

$$\mathcal{L}_{\text{scalar}}^{(2)(SD)} \sim \frac{1}{4\pi^2} \left(\frac{e^2}{4\pi} \right)^2 f^2 \ln \left(\frac{2ef}{m^2} \right) \quad (25)$$

From the coefficient of the $f^2 \ln f$ term we can read off the two-loop contribution to the scalar QED β -function, so that to two-loop order:

$$\beta_{\text{scalar}} = \frac{\alpha^2}{6\pi} + \frac{\alpha^3}{2\pi^2} + \dots \quad (26)$$

Once again, this illustrates the connection between strong-field and short-distance physics [8–10,31].

As a final application of our result, we consider the analytic continuation of $\mathcal{L}_{\text{scalar}}^{(2)(SD)}(\kappa)$ under $\kappa \rightarrow i\kappa$ (or, $f \rightarrow -if$). We obtain the following explicit Schwinger-type formula:

$$\text{Im}\mathcal{L}_{\text{scalar}}^{(2)(SD)}(i\kappa) = \alpha\pi \frac{m^4}{(4\pi)^3} \frac{1}{\kappa^2} \sum_{k=1}^{\infty} \left[2\pi\kappa k - 1 + 3\kappa^2 \text{Re} \left(\tilde{\psi}(i\kappa) \right) \right] \exp[-2\pi\kappa k] \quad (27)$$

where $\tilde{\psi}(x) \equiv \psi(x) - \ln(x) + \frac{1}{2x}$ is the same function that appears in the definition of $\xi(x)$ in Eq.(19). Note that the imaginary part (27) is non-perturbative in the field [recall Eq.(4)]. This is true also for the one-loop case:

$$\text{Im}\mathcal{L}_{\text{scalar}}^{(1)(SD)}(i\kappa) = \frac{m^4}{(4\pi)^3} \frac{1}{\kappa^2} \sum_{k=1}^{\infty} \left[\frac{2\pi\kappa}{k} + \frac{1}{k^2} \right] \exp[-2\pi\kappa k] \quad (28)$$

It is interesting that one finds a completely explicit expression for the exponential prefactor in (27), for any k . This should be contrasted with the two-loop constant electric field case where only a two-parameter integral representation is available [8–10], and where a complicated analysis [35] of the analyticity properties of the integrand led to an analogous representation for the imaginary part. But only the leading term in the κ expansion of the prefactor could be determined. A Borel analysis [15] of the high orders of perturbation theory, in this two-loop electric case, found a numerical estimate for the next term in this Ritus-Lebedev κ expansion of the prefactor (for $k = 1$), but was unable to go further. Yet, in the self-dual background studied here, we find a simple expression for the *entire* prefactor, for *all* k . We note that for a given value of κ , the two-loop prefactor in (27) eventually becomes larger than the corresponding one-loop prefactor, when the ‘instanton’ index k is large enough. If this were found to be true also in the electric case, one would have to conclude that the one-loop Schwinger formula (2) does not always provide a reliable estimate of the actual multi-pair production rate, for very large numbers of pairs.

We have also tested [23] the Borel dispersion relation approach on the self-dual case two-loop weak-field expansion (21), and find perfect agreement with the leading ($k = 1$) term in the exact result (27). Indeed, one can already see that the leading terms in (9) and (23) are consistent with the leading large κ terms in the imaginary parts (28) and (27).

To conclude, we reiterate that we find it extremely surprising that such a simple closed-form can be obtained for the two-loop effective Lagrangian. It suggests that even higher loop orders might be accessible in the self-dual case, and that the function $\xi(x)$ will reappear in them. This might ultimately lead to progress in the computation of higher beta function coefficients (only the first three coefficients are presently known for scalar QED [36]), as well as provide partial information on higher order light-by-light scattering [37]. Similar simplifications occur in the case of spinor QED [23]. A question of obvious interest is whether some of this simplification carries over also to the nonabelian case. Work in this direction is in progress.

Acknowledgements: We thank D. Broadhurst, H. Gies, M. Reuter and V. I. Ritus for helpful conversations and correspondence. CS thanks the Institute for Advanced Study, Princeton, for hospitality. GD thanks the US Dept. of Energy for support through grant DE-FG02-92ER40716.

- [1] W. Heisenberg and H. Euler, “Folgerungen aus der Diracschen Theorie des Positrons”, *Z. Phys.* **98** (1936) 714.
- [2] V. Weisskopf, “Über die Elektrodynamik des Vakuums auf Grund der Quantentheorie des Elektrons”, *Kong. Dans. Vid. Selsk. Math-fys. Medd.* XIV No. 6 (1936), reprinted in *Quantum Electrodynamics*, J. Schwinger (Ed.) (Dover, New York, 1958).
- [3] S. L. Adler, “Photon splitting and photon dispersion in a strong magnetic field”, *Ann. Phys. (N.Y.)* **67** (1971) 599.
- [4] J. Schwinger, “On gauge invariance and vacuum polarization”, *Phys. Rev.* **82** (1951) 664.
- [5] A. Nikishov, “Pair production by a constant external field”, *Zh. Eksp. Teor. Fiz.* **57** (1969) 1210 [*Sov. Phys. JETP* **30** (1970) 660].
- [6] D. L. Burke et al., “Positron production in multi-photon light-by-light scattering”, *Phys. Rev. Lett.* **79** (1997) 1626.
- [7] A. Ringwald, “Pair production from vacuum at the focus of an X-Ray free electron laser”, *Phys. Lett. B* **510** (2001) 107, (hep-ph/0103185).
- [8] V. I. Ritus, “Lagrangian of an intense electromagnetic field and quantum electrodynamics at short distances”, *Zh. Eksp. Teor. Fiz* **69** (1975) 1517 [*Sov. Phys. JETP* **42** (1975) 774].
- [9] V. I. Ritus, “Connection between strong-field quantum electrodynamics with short-distance quantum electrodynamics”, *Zh. Eksp. Teor. Fiz* **73** (1977) 807 [*Sov. Phys. JETP* **46** (1977) 423].
- [10] V. I. Ritus, “The Lagrangian Function of an Intense Electromagnetic Field”, in *Proc. Lebedev Phys. Inst. Vol. 168, Issues in Intense-field Quantum Electrodynamics*, V. I. Ginzburg, ed., (Nova Science Pub., NY 1987).
- [11] V. Fock, “Die Eigenzeit in der klassischen und in der Quantenmechanik”, *Phys. Z. Sowjet.* **12** (1937) 404.
- [12] W. Dittrich and M. Reuter, *Effective Lagrangians in Quantum Electrodynamics*, Springer 1985.
- [13] M. Reuter, M. G. Schmidt and C. Schubert, “Constant external fields in gauge theory and the spin 0, 1/2, 1 path integrals”, *Ann. Phys. (N.Y.)* **259** (1997) 313, (hep-th/9610191).
- [14] D. Fliegner, M. Reuter, M. G. Schmidt, C. Schubert, “The two-loop Euler-Heisenberg Lagrangian in dimensional renormalization”, *Theor. Mat. Fiz.* **113** (1997) 289 [*Theor. Math. Phys.* **113** (1997) 1442]; C. Schubert, “Perturbative quantum field theory in the string-inspired formalism”, *Phys. Rept.* **355** (2001) 73, (hep-th/0101036).
- [15] G. V. Dunne and C. Schubert, “Two-loop Euler-Heisenberg QED pair-production rate”, *Nucl. Phys. B* **564** (2000) 591, (hep-th/9907190).
- [16] Z. Bern and D. A. Kosower, “Efficient calculation of one loop QCD amplitudes”, *Phys. Rev. Lett.* **66** (1991) 1669, “The computation of loop amplitudes in gauge theory”, *Nucl. Phys. B* **379** (1992) 451.
- [17] M. J. Strassler, “Field theory without Feynman diagrams: one-loop effective actions”, *Nucl. Phys. B* **385** (1992) 145, (hep-ph/9205205).
- [18] D.G.C. McKeon and T.N. Sherry, “Radiative effects in a constant magnetic field using the quantum mechanical path integral”, *Mod. Phys. Lett. A* **9** (1994) 2167.
- [19] M. G. Schmidt and C. Schubert, “Multiloop calculations in the string-inspired formalism: the single spinor-loop in QED”, *Phys. Rev. D* **53** (1996) 2150, (hep-th/9410100).
- [20] R. Zh. Shaisultanov, “On the string-inspired approach to QED in external field”, *Phys. Lett. B* **378** (1996) 354, (hep-th/9512142).
- [21] C. Schubert, “Perturbative quantum field theory in the string-inspired formalism”, *Phys. Rept.* **355** (2001) 73, (hep-th/0101036).
- [22] B. Körs and M.G. Schmidt, “The effective two loop Euler-Heisenberg action for scalar and spinor QED in a general constant background field”, *Eur. Phys. J. C* **6** (1999) 175, (hep-th/9803144).
- [23] G. V. Dunne and C. Schubert, in preparation.
- [24] H. Leutwyler, “Vacuum fluctuations surrounding soft gluon fields”, *Phys. Lett. B* **96** (1980) 154, “Constant gauge fields and their quantum fluctuations”, *Nucl. Phys. B* **179** (1981) 129.
- [25] O-K. Kwon, C. Lee and H. Min, “Massive field contributions to the QCD vacuum tunneling amplitude”, *Phys. Rev. D* **62** (2000) 114022, (hep-ph/0008028).
- [26] H. Liu and A.A. Tseytlin, “D3-brane - D-instanton configuration and N=4 super YM theory in constant self-dual background”, *Nucl. Phys. B* **553** (1999) 231, (hep-th/9903091).
- [27] I.L. Buchbinder, A.Yu. Petrov and A.A. Tseytlin, “Two-loop N = 4 Super Yang Mills effective action and interaction between D3-branes”, *OHSTPY-HEP-T-01-22* (hep-th/0110173).
- [28] G. V. Efimov, A. C. Kalloniatis and S. N. Nedelko, “Confining Properties of the Homogeneous Self-Dual Field and the Effective Potential in SU(2) Yang-Mills Theory”, *Phys. Rev. D* **59** (1999) 014026, (hep-th/9806165).
- [29] S. Blau, M. Visser and A. Wipf, “Analytic results for the effective action”, *Int. J. Mod. Phys. A* **6** (1991) 5409.
- [30] This function is closely related to Barnes’s G-function: E. W. Barnes, “The theory of the G-function”, *Quart. J. Pure Appl. Math* **XXXI** (1900) 264; E. T. Whittaker and G. N. Watson, *Modern Analysis*, (Cambridge University Press, 1927)

page 264.

- [31] S. G. Matinyan and G. K. Savvidy, “Vacuum polarization induced by the intense gauge field”, Nucl. Phys. B **134** (1978) 539.
- [32] C. Itzykson and J. Zuber, *Quantum Field Theory*, (McGraw-Hill, New York, 1985).
- [33] C. Schubert, “Vacuum polarisation tensors in constant electromagnetic fields: Part I”, Nucl. Phys. B **585** (2000) 407, (hep-ph/0001288).
- [34] J. Iliopoulos, C. Itzykson and A. Martin, “Functional methods and perturbation theory”, Rev. Mod. Phys. **47** (1975) 165.
- [35] S. L. Lebedev and V. I. Ritus, “Virial representation of the imaginary part of the Lagrangian function of an electromagnetic field”, Zh. Eksp. Teor. Fiz. **86** (1984) 408 [JETP **59** (1984) 237].
- [36] D.J. Broadhurst, R. Delbourgo and D. Kreimer, “Unknotting the polarized vacuum of quenched QED”, Phys. Lett. B **366** (1996) 421, (hep-ph/9509296).
- [37] Z. Bern, A. De Freitas, L. Dixon, A. Ghinculov and H. L. Wong, “QCD and QED corrections to light-by-light scattering”, (hep-ph/0109079).